

Assessment of Self-Medication Use among University Students

*Abdullah Alkhalwaldeh¹, PhD, Omar Al Omari², PhD, Mohammed ALBashtawy³, PhD, Omar Khraisat⁴, PhD, Khloud Al Dammery⁵, MSc, Shadi F. Gharaibeh⁶, PhD, Ibrahim Ayasrah,⁷ MSc

Abstract

Introduction: The use of self-medication represents a health problem among university students. **Aim:** This study aimed to explore the self-medication phenomenon among university students. **Methods:** Using a cross-sectional design and self-reported questionnaire, data were collected from 150 university students.

Results: Use of self-medication was reported by 69.3% of the students. The most frequent medications used were analgesics (61.3%), cold & cough medications (58%) and antibiotics (56.7%), with the most common health conditions for self-medication being headaches (64.7%), colds and flu (62.7%), and fever (52.0%). The ease of access to medication (36.0%), advice from friends (32.7%), lack of time to consult physicians (27.3%), long waiting times for treatment (26.7%), hospital medications do not work (20.0%), and the hospital or clinic is very far away (18.7%) were reported by the students as reasons leading to self-medication. The most common sources for self-medication were pharmacies (58.7%), and available in home (49.3%), and hospital (31.3%). **Conclusion:** The use of self-medication is high among university students. Therefore, decision makers should pay more attention to this phenomenon and more education sessions should be given to youth to draw their attention to the possible negative effects of self-medication.

Keywords: Self-medication; Use; University Students; Jordan.

1. Introduction

The use of self-medication is a global health issue (Sarahroodi, Maleki-Jamshid, Sawalha, Mikaili, & Safaiean, 2012), and it represents a health problem among universities students (Klemenc-Ketis, Hladnik, & Kersnik, 2010). Its described in the literature (Helal & Abou-ElWafa, 2017; Al-Ameri, Al-Badri & Lafta, 2017; Badiger et al., 2012). A cross-sectional study of 800 university student found that approximately 62.9% of the students used self-medication (Helal & Abou-ElWafa, 2017). Approximately, 92% of the university students used self-medication in India (Badiger et al., 2012), and in Iraq (Al-Ameri et al., 2017).

The reasons for using of self -medications by university students include: the belief that health problems were mild and did not deserve a physician consultation (Gama & Secoli, 2017; Albusalih, Naqvi, Ahmad & Ahmad, 2017;; Sawhney et al., 2017). Lack of time to consult a physician (Gama & Secoli, 2017; Shehnaz, Khan & Sreedharan, 2013) and prior experience with medications (Albusalih, et al., 2017; Shehnaz, Khan, & Sreedharan, 2013) were another reasons also. Furthermore, advice from friend and to avoid cost of consultation (Shehnaz, Khan, & Sreedharan, 2013). Lastly, ease of access to medications (Syed et al., 2014) and to avoid long waiting time (Gutema et al., 2011).

¹ Associate Professor, Faculty of Nursing, Jerash University, Jerash, Jordan. Email: dr-abd@jpu.edu.jo.

² Associate Professor, Faculty of Nursing, Sultan Qaboos University, Muscat, Oman. Email: o.alomari@squ.edu.om

³ Professor, Princess Salma Faculty of Nursing, AL al-Bayt University, Mafraq, Jordan Email: mohammadbash@aabu.edu.jo

⁴ Assistant Professor, Faculty of Nursing, Al-Ahliyya Amman University, Amman, Jordan. Email: khraisat1111@hotmail.com

⁵ Faculty of Nursing, Sultan Qaboos University, Muscat, Oman. Email: kdameery@yahoo.com

⁶ Associate Professor, Faculty of Pharmacy, Jerash University, Jerash, Jordan Email: shadi.gharaibeh@jpu.edu.jo

⁷ Faculty of Nursing, Jerash University, Jerash, Jordan. Email: ibrahim.ayasrah@jpu.edu.jo

The main sources for self-medication among university students include: the pharmacy (Bunduki, Mumbere, & Mbahweka, 2017; Rock Britto, Elango, Shilpa Charles, Shyamala Priya, & Sivaranjan, 2017); friends and relatives (Rock Britto et al., 2017; Ahmadi et al., 2016; Gutema et al., 2011), private clinic and hospital (Bunduki et al., 2017). A number of health conditions were found to promote students' use of self-medication; these were headache, cold and fever (Ahmadi, Jamshidi, Sadeghi, Abdi, & Vahid, 2016; Syed, Naseer, Memon, & Rani, 2014; Saeed et al., 2014; Badiger et al., 2012). In Iran, other health conditions such as allergy, insufficient vitamins and minerals, respiratory disease, and gastrointestinal problems were reported (Zardosht et al., 2016). While, in Pakistan, storage of medications for multi-purposes, occasional pain, common infections, and cough/cold were the health conditions that promote use of self-medication (Ullah et al., 2013).

Previous studies revealed that the analgesics, antibiotics, and antipyretics were the most commonly used self-medications among university students (Al-Ameri et al., 2017; Sawhney, Bhat, & Singh, 2017). A cross-sectional study from Iran found that cold and cough medications were the most commonly used medications followed by analgesics, antihistamines, herbal drugs, vitamins, dietary supplements, minerals, energizers, and antibiotics (Zardosht et al., 2016).

In Jordan, previous studies focused on the assessment of self-medication among public schools' students (ALBashtawy, Batiha, Tawalbeh, Tubaishat, & AlAzzam, 2015), and among students of medical faculties in public universities (Alkhatabeh, Alefan, & Alqudah, 2016). The current study explored self-medication phenomena among university students. That is, it assessed the prevalence self-medication among Jordanian universities students. Health conditions that promoted universities students to use self-medication was also explored and lastly main reasons and source for self-medication were investigated.

2. Methods

2.1 Design

A cross sectional study was conducted to assess the prevalence of self medication, the type of medication students use, health conditions that promote self medication use, and source of self medications among the students from September to October 2019.

2.2 Participants

A convenient sample of 150 universities students participated in the current study. Inclusion criteria were:

- Student registered for the first semester 2019-2020.
- Student has Jordanian nationality.

The exclusion criteria were being an international student.

2.3 Instrument

The questionnaire, which was based on the literature review, consisted of three parts: Socio-demographic: This part was used to gather data on each student's age (years), gender (male or female), faculty (medical; nursing & pharmacy) and (non medical; humanities, economic & accounting), level of academic year (first, second, third, fourth or more).

Use of self-medication: This part was assessed retrospectively by asking students the following: Have you used medications by yourself (without prescription and without seeking medical consultation) in the past 6 months? For positive responses, students were then asked, What medications have you used? And, what is the purpose (health conditions/disease) of taking these medications?

Reasons for self-medication and medication source: Reasons behind self-medication were assessed by asking students the following: What are the reasons that force you to use medications by yourself (without medical prescription)?

Medication source was assessed by asking students the following: What are the sources of self-medication?

2.4 Ethical Considerations

The study was approved by ethics committee of the University. Preceding data collection, the aim of the study was clarified to students, and informed consent was obtained. Also, anonymity and confidentiality of the students were maintained.

2.5 Statistical Analysis

Descriptive statistics (frequencies, mean and percentages) were applied to analyze data using Statistical Package for Social Sciences (SPSS) version 17.

3. Results

3.1 Characteristics of students

Of the 150 students that participated in the present study, 81 (54.0%) were females. The mean age of participants was 24.93 years old (SD = 5.894). The faculties were classified into: medical faculties 68 (45.3%), and non medical 82 (54.7%). The dominant group among the students was the fourth year & more students 66 (44.0%), whereas the least represented group was first year students 18 (12.0%) (see Table 1).

TABLE 1 Characteristics of students (N = 150)

| Variables | Frequency | Percentage (%) | Mean (S.D) |
|-------------------------------|-----------|----------------|---------------|
| Age | | | 24.93 (5.894) |
| Gender: | | | |
| Male | 69 | 46.0 | |
| Female | 81 | 54.0 | |
| Faculty: | | | |
| Medical faculties | 68 | 45.3 | |
| Nonmedical faculties | 82 | 54.7 | |
| level of academic year | | | |
| First year | 18 | 12.0 | |
| Second year | 27 | 18.0 | |
| Third year | 39 | 26.0 | |
| Fourth year & more | 66 | 44.0 | |

3.2 Use of self-medication

About 104 (69.3%) of the students had used the self-medication at least once in the past 6 months and about 46 (30.7%) had not used the self-medication. Among the students who used the self-medication in the past 6 months, the most frequent medications for use were analgesics 92(61.3%), cold & cough medications 87(58.0%) and antibiotics 85(56.7%), while the less frequent medications for use were antispasmodic 5(3.3%), stimulants 5(3.3%) and psychotropic medications 1 (0.7%) (see Table 2).

Table 2 Use of Self-Medication (N = 150)

| | Frequency | Percentage (%) |
|---------------------------------------------------|-----------|----------------|
| Use of self-medication | | |
| Yes | 104 | 69.3 |
| No | 46 | 30.7 |
| Medication used: | | |
| Analgesics | 92 | 61.3 |
| Cold & cough medicines | 87 | 58.0 |
| Antibiotics | 85 | 56.7 |
| Antipyretics | 80 | 53.3 |
| Herbals | 73 | 48.7 |
| Antacids | 47 | 31.3 |
| Vitamins & nutritional complements | 43 | 28.7 |
| Medications for treatment of hair & skin diseases | 39 | 26.0 |
| Applied Creams | 31 | 20.7 |
| Aspirin | 24 | 16.0 |
| Anti- emetics | 20 | 13.3 |
| Antihistamines | 19 | 12.7 |
| Anti-diarrheal | 17 | 11.3 |
| Laxatives | 15 | 10.0 |
| Antidepressants and Anti-anxiety | 9 | 6.0 |
| Cardiovascular medications | 7 | 4.7 |
| Antispasmodic | 5 | 3.3 |
| Stimulants | 5 | 3.3 |
| Psychotropic medications | 1 | 0.7 |

3.3 Health conditions for self-medication

The most common health conditions behind self-medication were headache 97(64.7%), cold and flu 94 (62.7%), fever 78(52.0%), while the less frequent health conditions for self-medication use were Psychological problems 14(9.3%), Skin diseases 14(9.3%), and Bacterial & viral infections 14(9.3%) (see Table 3).

TABLE 3 Health conditions that make students use self-medication (N = 150)

| | Frequency | Percentage (%) |
|-----------------------------------------------|-----------|----------------|
| Health conditions for self-medication: | | |
| Headache | 97 | 64.7 |
| Cold and flu | 94 | 62.7 |
| Fever | 78 | 52.0 |
| Toothache | 66 | 44.0 |
| Allergy | 41 | 27.3 |
| Nausea/vomiting | 39 | 26.0 |
| Constipation | 34 | 22.7 |
| Inability to sleep | 34 | 22.7 |
| Acid reflux | 30 | 20.0 |
| Muscle Spasm | 29 | 19.3 |
| Diarrhea | 28 | 18.7 |
| Respiratory problems | 26 | 17.3 |
| Eye Problems | 25 | 16.7 |
| Weight loss or gain purpose | 23 | 15.3 |
| Dysmanohrea | 20 | 13.3 |
| Genitourinary infections | 17 | 11.3 |
| Psychological problems | 14 | 9.3 |
| Skin diseases | 14 | 9.3 |
| Bacterial & viral infections | 14 | 9.3 |

3.4 Students's Perception of Reasons Affecting Their Use of Self-Medication

Table 4 shows that the main reasons behind use of self-medication were related to ease of access to medication 54(36.0%), advice from friend 49 (32.7%), lack of time to consult physician 41(27.3%), long waiting times for treatment 40 (26.7%), hospital medications do not work 30(20.0%), the hospital or clinic is very far away 28 (18.7%). Cost of consultation 26(17.3%) was the reason least perceived by students to affect their use of self-medication.

TABLE 4 Reasons for self-medication (N = 150)

| Reason | Frequency | Percentage (%) |
|-----------------------------------------|-----------|----------------|
| Ease of access to medications | 54 | 36.0 |
| Advice from friend | 49 | 32.7 |
| Lack of time to consult physician | 41 | 27.3 |
| Long waiting time | 40 | 26.7 |
| Hospital medications do not work | 30 | 20.0 |
| The hospital or clinic is very far away | 28 | 18.7 |
| Cost of consultation | 26 | 17.3 |

3.5 Source of self-medication

The most common source for self-medication was pharmacy 88(58.7%), while the less frequent source for self-medication was friends 31(20.7%) (see Table 5).

TABLE 5 Source of self-medication (N = 150)

| Source | Frequency | Percentage (%) |
|-----------------------|-----------|----------------|
| Pharmacy | 88 | 58.7 |
| Available in home | 74 | 49.3 |
| Hospital | 47 | 31.3 |
| Health staff | 47 | 31.3 |
| Previous prescription | 43 | 28.7 |
| Friends | 31 | 20.7 |

4. Discussion

In the present study, the rate of self-medication among university students in the past 6 months was 69.3%, and this is close to the rate reported in previous literature (Helal & Abou-ElWafa, 2017) but, less than the prevalence that was found among university students in other countries (Al-Ameri et al., 2017; Badiger et al., 2012). One explanation for such difference could be attributed to the discrepancy in demographic characteristics, socioeconomic status, and because of differences in methodologies and recall periods used to assess the use of self-medication among university students. there is a need to increase students awareness on appropriate use of self-medications because of many risks of self-medication (e.g.: medication adverse effects and dependency, medication and food interaction, and resistance of microorganisms) (WHO, 2000).

The current study found that analgesics, cold & cough medications and antibiotics were the most frequent self-medications used among the universities students, which is in line with previous international studies (Al-Ameri et al., 2017; Zardosht et al., 2016; Sawhney et al., 2017). The irresponsible or/and overuse of analgesics self-medication is associated with side effects (Sarahroodi et al., 2012). For example, misuse of Non Steroidal Anti inflammatory Drugs (NSAIDs) may cause renal failure and hepatic dysfunction (Bashrahlil & Baruzraig, 2008). As well, the use of antibiotics self-medication contributes in the development of resistance of microorganisms to antibiotics (Morgan, Okeke, Laxminarayan, Perencevich, & Weisenberg, 2011). There is a need to increase university students' awareness regarding complications and contraindications of these medications in order to avoid the complication of using it. Future research needed to assess the use of analgesics and antibiotics self-medication in details among the university students in Jordan.

In the current study, the main health conditions behind using self-medication were headache, cold and flu, and fever. This is similar to many other prior studies (Ahmadi et al., 2016; Syed et al., 2014; Saeed et al., 2014; Badiger et al., 2012).

University students in the current study and previous studies used self-medication because the ease of access to medication (Ahmadi et al., 2016), and advice from friend. Cost of consultation (Shehnaz et al., 2013), lack of time to consult physician (Gama & Secoli, 2017; Shehnaz et al., 2013), and long waiting times for treatment (Gutema et al., 2011) were another motivating factors.

The current study found that pharmacy was the main source for self-medication. Other frequent sources were available in home, hospital, health staff, previous prescription and friends. These findings were close to other recent study (Bunduki et al., 2017; Rock Britto et al., 2017; Ahmadi et al., 2016; Zardosht et al., 2016). Consequently, the pharmacists role is important by selling medications only with medications prescriptions.

4.1 Limitations of the Study

Use of self-medication was self-reported which may result in recall bias. Nevertheless, it was measured as a dichotomous variable, which is easier to recall.

4.2 Conclusion and Recommendations

The findings reflected high rates of use of self-medication among Jordanian universities students'. This study added to the body of knowledge in nursing regarding health-seeking behavior of the university students, which will help in promoting university students' health. Future plans to improve universities students health must address the reasons that force student to use self-medication. Solutions to address this problem might consist of (a) offer health education programs to raise awareness of the university students regarding the possible problems of self-medication use to reduce risks; and (b) enforce medication laws and manage the pharmacies medications sale.

Conflict of Interest

No conflict of interests concerning the publication of the present study.

References

- Ahmadi, S. M., Jamshidi, K., Sadeghi, K., Abdi, A., & Vahid, M. P. (2016). The prevalence and affecting factors on self-medication among students of Kermanshah University of Medical Science in 2014. *Journal of clinical and diagnostic research: JCDR*, 10(5), IC01.
- Al-Ameri, R. J., Al-Badri, H. J. A., & Lafta, R. K. (2017). Prevalence of self-medication among university students in Baghdad: a cross-sectional study from Iraq/Prévalence de l'automédication parmi les étudiants à l'Université de Bagdad: étude transversale iraquienne. *Eastern Mediterranean Health Journal*, 23(2), 87.
- ALBashtawy, M., Batiha, A. M., Tawalbeh, L., Tubaihsat, A., & AlAzzam, M. (2015). Self-medication among school students. *The Journal of School Nursing*, 31(2), 110-116. <https://doi.org/10.1177/1059840514554837>
- Albusalih, F. A., Naqvi, A. A., Ahmad, R., & Ahmad, N. (2017). Prevalence of self-medication among students of pharmacy and medicine colleges of a public sector university in Dammam City, Saudi Arabia. *Pharmacy*, 5(3), 51. <https://doi.org/10.3390/pharmacy5030051>
- Alkhatatbeh, M. J., Alefan, Q., & Alqudah, M. A. (2016). High prevalence of self-medication practices among medical and pharmacy students: a study from Jordan. *Int J Clin Pharmacol Ther*, 54(5), 390-8. <https://doi.org/10.5414/CP202451>.
- Badiger, S., Kundapur, R., Jain, A., Kumar, A., Pattanshetty, S., Thakolkaran, N., ... & Ullal, N. (2012). Self-medication patterns among medical students in South India. *The Australasian medical journal*, 5(4), 217. <https://doi.org/10.4066/AMJ.2012.1007>
- Bashrahil, K. A., & Baruzaiq, A. S. (2008). Self-Medication: Concept, Prevalence & risks in Mukalla City (Yemen) 2004-2005. *Andalus for studies & Research*, 2, 1-3.
- Bunduki, G. K., Mumbere, M., & Mbahweka, F. K. (2017). Assessment of antibiotic self-medication pattern among university students in Eastern Democratic Republic of the Congo. *Journal of Pharmaceutical Research International*, 18(1), 1-7. <https://doi.org/10.9734/JPRI/2017/31848>
- Gama, A. S. M., & Secoli, S. R. (2017). Self-medication among nursing students in the state of Amazonas–Brazil. *Revista gaucha de enfermagem*, 38(1). <https://doi.org/10.1590/1983-1447.2017.01.65111>.
- Gutema, G. B., Gadisa, D. A., Kidanemariam, Z. A., Berhe, D. F., Berhe, A. H., Hadera, M. G., ... & Abrha, N. G. (2011). Self-medication practices among health sciences students: the case of Mekelle University. *Journal of Applied Pharmaceutical Science*, 1(10), 183.

- Helal, R. M., & Abou-ElWafa, H. S. (2017). Self-Medication in University Students from the City of Mansoura, Egypt. *Journal of environmental and public health*, 2017. <https://doi.org/10.1155/2017/9145193>
- Klemenc-Ketis, Z., Hladnik, Z., & Kersnik, J. (2010). Self-medication among healthcare and non-healthcare students at University of Ljubljana, Slovenia. *Medical Principles and practice*, 19(5), 395-401. <https://doi.org/10.1159/000316380>
- Morgan, D. J., Okeke, I. N., Laxminarayan, R., Perencevich, E. N., & Weisenberg, S. (2011). Non-prescription antimicrobial use worldwide: a systematic review. *The Lancet infectious diseases*, 11(9), 692-701. [https://doi.org/10.1016/S1473-3099\(11\)70054-8](https://doi.org/10.1016/S1473-3099(11)70054-8)
- Rock Britto, R. S., Elango, S., Shilpa Charles, D., Shyamala Priya, M., & Sivarajan, R. (2017). Prevalence and Pattern of Self-Medication among Undergraduate Medical Students. *International Journal of Innovative Research in Medical Science*, 2(06), 862-to.
- Saeed, M. S., Alkhalhaiban, A. S., Al-Worafi, Y. M. A., & Long, C. M. (2014). Perception of self-medication among university students in Saudi Arabia. *Archives of Pharmacy Practice*, 5(4), 149. <https://doi.org/10.4103/2045-080X.142049>
- Sarahroodi, S., Maleki-Jamshid, A., Sawalha, A. F., Mikaili, P., & Safaeian, L. (2012). Pattern of self-medication with analgesics among Iranian University students in central Iran. *Journal of family & community medicine*, 19(2), 125-9. <https://doi.org/10.4103/2230-8229.98302>
- Sawhney, V., Bhat, M. Y., & Singh, Z. (2017). A descriptive study of prevalence, pattern and attitude of self-medication among second professional medical students in a tertiary care center. *International Journal of Basic & Clinical Pharmacology*, 4(3), 542-546. <https://doi.org/10.18203/2319-2003>.
- Shehnaz, S. I., Khan, N., & Sreedharan, J. Prevalence and practice of self-medication among medical students. *GULF MEDICAL JOURNAL*. 2013;2(S2):S86-S92
- Syed, N., Naseer, M., Memon, M. Q., & Rani, K. (2014). Prevalence of self-medication and its practice among the medical and non-medical students. *JLUMHS*, 13(02), 79.
- Ullah, H., Khan, S. A., Ali, S., Karim, S., Baseer, A., Chohan, O., ... & Murtaza, G. (2013). Evaluation of self-medication amongst university students in Abbottabad, Pakistan; prevalence, attitude and causes. *Acta Pol Pharm*, 70(5), 919-22.
- World Health Organization. (2000). Guidelines for the regulatory assessment of medicinal products for use in self-medication.
- Zardosht, M., Dastoorpoor, M., Hashemi, F. B., Estebarsari, F., Jamshidi, E., Abbasi-Ghahramanloo, A., & Khazaeli, P. (2016). Prevalence and Causes of Self Medication among Medical Students of Kerman University of Medical Sciences, Kerman, Iran. *Global journal of health science*, 8(11), 150.